

A summary of influenza surveillance indicators reported to MDH for the week ending March 9, 2019

Prepared by the Division of Infectious Disease Surveillance Prevention and Health Promotion Administration Maryland Department of Health

The data presented in this document are provisional and subject to change as additional reports are received.

SUMMARY

During the week ending March 9, 2019 influenza-like illness (ILI) intensity in Maryland was MODERATE and there was WIDESPREAD geographic activity. The proportion of outpatient visits for ILI reported by Sentinel Providers and by Maryland Emergency Departments increased. The proportion of MRITS respondents reporting ILI increased from last week. Clinical laboratories reported a slight increase in the proportion of specimens testing positive for influenza. Two hundred and forty four specimens tested positive for influenza at the MDH lab. There were 196 influenza-associated hospitalizations. There were ten respiratory outbreaks reported to MDH.


Click here to visit our influenza surveillance web page

ILI Intensity Levels			
Minimal			
Low			
✓ Moderate			
High			

Influenza Geographic Activity
No Activity
Sporadic
Local
Regional
√Widespread

ILINet Sentinel Providers


Nineteen providers reported a total of 6,252 visits this week. Of those, 251 (4.0%) were visits for ILI. This is above the Maryland baseline of 2.0%.



ILI Visits To Sentinel Providers By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	80 (32%)	119 (44%)	1,183 (32%)
Age 5-24	134 (53%)	107 (39%)	1,495 (41%)
Age 25-49	17 (7%)	24 (9%)	534 (14%)
Age 50-64	14 (6%)	12 (4%)	292 (8%)
$Age \ge 65$	6 (2%)	11 (4%)	179 (5%)
Total	251 (100%)	273 (100%)	3,683 (100%)

Visits to Emergency Departments for ILI

Emergency Departments in Maryland reported a total of 61,029 visits this week through the ESSENCE surveillance system. Of those, 4,146 (6.8%) were visits for ILI.



ILI Visits To Emergency Departments By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	639 (15%)	582 (15%)	9,347 (20%)
Age 5-24	1,326 (32%)	1,127 (30%)	13,873 (30%)
Age 25-49	1,381 (33%)	1,292 (34%)	14,378 (31%)
Age 50-64	533 (13%)	517 (14%)	5,847 (13%)
Age ≥ 65	267 (6%)	267 (7%)	3,035 (7%)
Total	4,146 (100%)	3,785 (100%)	46,480 (100%)

Neighboring states' influenza information:

Delaware http://dhss.delaware.gov/dph/epi/influenzahome.html

District of Columbia http://doh.dc.gov/service/influenza


Pennsylvania http://www.health.pa.gov/My%20Health/Diseases%20and%20Conditions/I-L/Pages/Influenza.aspx#.V-LtaPkrJD8

 $\begin{tabular}{ll} Virginia & \underline{http://www.vdh.virginia.gov/epidemiology/influenza-flu-in-virginia/influenza-surveillance/} \\ \end{tabular}$

West Virginia http://dhhr.wv.gov/oeps/disease/flu/Pages/fluSurveillance.aspx

A summary of influenza surveillance indicators reported to MDH for the week ending March 9, 2019


MRITS is the Maryland Resident Influenza Tracking System, a weekly survey for influenza-like illness (ILI). A total of 574 residents responded to the MRITS survey this week. Of those, 17 (3.0%) reported having ILI and missing 26 cumulative days of regular daily activities.




MRITS Respondents Reporting ILI By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	1 (6%)	0 (0%)	13 (7%)
Age 5-24	6 (35%)	4 (36%)	47 (26%)
Age 25-49	3 (18%)	1 (9%)	50 (28%)
Age 50-64	7 (41%)	3 (27%)	39 (22%)
Age ≥ 65	0 (0%)	3 (27%)	29 (16%)
Total	17 (100%)	11 (100%)	178 (100%)

Clinical Laboratory Influenza Testing


There were 63 clinical laboratories reporting 8,396 influenza diagnostic tests, mostly rapid influenza diagnostic tests (RIDTs). Of those, 2,371 (28.2%) were positive for influenza. Of those testing positive, 2,181 (92%) were influenza Type A and 190 (8%) were influenza Type B. The reliability of RIDTs depends largely on the conditions under which they are used. False-positive (and true-negative) results are more likely to occur when the disease prevalence in the community is low, which is generally at the beginning and end of the influenza season and during the summer.





State Laboratories Administration Influenza Testing

The MDH Laboratories Administration performed a total of 315 PCR tests for influenza and 244 (77.5%) were positive for influenza. Of those testing positive, 159 (65%) were positive for Type A (H1), 81 (33%) were positive for Type A (H3), and 4 (2%) was positive for Type B (Yamagata). PCR testing is more reliable than RIDT. The MDH testing identifies subtypes of influenza A and lineages of influenza B, information that is not available from the RIDT results. The table below summarizes results by type, subtype, and lineage.




Positive PCR Tests by Type (Subtype)	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A (H1)	159 (65%)	381 (75%)	1,564 (75%)
Type A (H3)	81 (33%)	125 (25%)	442 (21%)
Type B (Victoria)	0 (0%)	3 (1%)	62 (3%)
Type B (Yamagata)	4 (2%)	1 (<1%)	12 (1%)
Dual Type A (H1/H3)	0 (0%)	0 (0%)	1 (<1%)
Total	244 (100%)	510 (100%)	2,081 (100%)

A summary of influenza surveillance indicators reported to MDH for the week ending March 9, 2019

Influenza-associated Hospitalizations

A total of 196 influenza-associated hospitalizations were reported this week. (A person with an overnight hospital stay along with a positive influenza test of any kind, e.g., RIDT or PCR, is considered an "influenza-associated hospitalization" for purposes of influenza surveillance.) This surveillance is conducted as a component of the Maryland Emerging Infections Program.



Influenza- Associated Hospitalizations by Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	16 (8%)	23 (9%)	166 (9%)
Age 5-17	9 (5%)	15 (6%)	99 (5%)
Age 18-24	2 (1%)	7 (3%)	38 (2%)
Age 25-49	32 (16%)	40 (15%)	327 (18%)
Age 50-64	50 (26%)	73 (28%)	522 (28%)
Age ≥ 65	87 (44%)	101 (39%)	710 (38%)
Total	196 (100%)	259 (100%)	1,862 (100%)


Influenza-associated Deaths

An influenza-associated death is one with a clinically compatible illness and a positive influenza test of any kind.

Pediatric Deaths: The total number of pediatric (< 18 years of age) deaths reported this influenza season is 1.

Influenza-associated pediatric mortality is a reportable condition in Maryland. Pediatric deaths are tracked without regard to hospitalization.


Adult Deaths Among Hospitalized Patients: A cumulative season total of 31 deaths have been reported among adults admitted to Maryland hospitals. Influenza-associated adult mortality is not a reportable condition in Maryland. However, surveillance for mortality in hospitalized adults is conducted as a



Influenza-Associated Deaths	Cumulative Season Total
Pediatric Deaths (Age < 18)	1
Adult Deaths (in hospitalized cases)	31

Outbreaks of Respiratory Disease

There were ten respiratory outbreaks reported to MDH this week. (Disease outbreaks of any kind are reportable in Maryland. Respiratory outbreaks may be reclassified once a causative agent is detected, e.g., from ILI to influenza.)




Respiratory Outbreaks by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Influenza	10 (100%)	13 (87%)	54 (68%)
Influenza-like Illness	0 (0%)	2 (13%)	14 (18%)
Pneumonia	0 (0%)	0 (0%)	12 (15%)
Other Respiratory	0 (0%)	0 (0%)	0 (0%)
Total	10 (100%)	15 (100%)	80 (100%)


A summary of influenza surveillance indicators reported to MDH for the week ending March 9, 2019

National Influenza Surveillance (CDC)

Influenza activity decreased slightly, but remains elevated in the United States. Influenza A(H1N1)pdm09, influenza A(H3N2), and influenza B viruses continue to co-circulate.


- Viral Surveillance: The percentage of respiratory specimens testing positive for influenza viruses in clinical laboratories decreased slightly. Nationally, during week 10, influenza A(H3) viruses were reported more frequently than influenza A(H1N1)pdm09 viruses. During the most recent three weeks, influenza A(H3) viruses were reported more frequently than influenza A(H1N1)pdm09 viruses in HHS Regions 2, 4, 5, 6, 7, 8 and 10.
- Influenza-like Illness Surveillance: The proportion of outpatient visits for influenza-like illness (ILI) decreased slightly to 4.5%, which is above the
 national baseline of 2.2%. All 10 regions reported ILI at or above their region-specific baseline level.
- Geographic Spread of Influenza: The geographic spread of influenza in Puerto Rico and 46 states was reported as widespread; four states reported regional
 activity; the District of Columbia reported local activity; the U.S. Virgin Islands reported sporadic activity; and Guam did not report.
- Influenza-associated Hospitalizations: A cumulative rate of 41.3 laboratory-confirmed influenza-associated hospitalizations per 100,000 population was reported. The highest hospitalization rate is among adults 65 years and older (123.9 hospitalizations per 100,000 population).
- Pneumonia and Influenza Mortality: The proportion of deaths attributed to pneumonia and influenza (P&I) was below the system-specific epidemic
 threshold in the National Center for Health Statistics (NCHS) Mortality Surveillance System.
- Influenza-associated Pediatric Deaths: Four influenza-associated pediatric deaths were reported to CDC during week 10.
- Outpatient Illness Surveillance: Nationwide during week 10, 4.5% of patient visits reported through the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) were due to influenza-like illness (ILI). This percentage is above the national baseline of 2.2%. (ILI is defined as fever (temperature of 100°F [37.8°C] or greater) and cough and/or sore throat.)





Data collected in ILINet may disproportionally represent certain populations within a state, and therefore, may not accurately depict the full picture of influenza activity for the whole state.

Data displayed in this map are based on data collected in ILINet, whereas the State and Territorial flu activity map is based on reports from state and territorial epidemiologists. The data presented in this map is preliminary and may change as more data are received. Differences in the data presented here by CDC and independently by some state health departments likely represent differing levels of data completeness with data presented by the state likely being the more complete.



Where to get an influenza vaccination

Interested in getting a flu vaccine for the 2018-19 influenza season? Go to https://phpa.health.maryland.gov/influenza/Pages/getvaccinated.aspx and click on your county/city of residence. You will be redirected to your local health department website for local information on where to get your flu vaccine.